



## Bifurcation Analysis of Structures in a Convection Model

**Dam, Magnus; Brøns, Morten; Rasmussen, Jens Juul; Naulin, Volker**

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Dam, M., Brøns, M., Rasmussen, J. J., & Naulin, V. (2015). *Bifurcation Analysis of Structures in a Convection Model*. Poster session presented at XXXV Dynamics Days Europe 2015, Exeter, United Kingdom.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Bifurcation Analysis of Structures in a Convection Model

Magnus Dam<sup>1</sup>, Morten Brøns<sup>1</sup>, Jens Juul Rasmussen<sup>2</sup>, and Volker Naulin<sup>2</sup>

<sup>1</sup>DTU Compute, Technical University of Denmark, Kgs. Lyngby, Denmark; <sup>2</sup>DTU Physics, Technical University of Denmark, Kgs. Lyngby, Denmark. Email: magnusd@dtu.dk

**This study investigates the convection driven evolution of isolated structures in a magnetized plasma. The analysis is based on numerical simulations of the evolution of two-dimensional plasma filaments in a nonuniform magnetic field. A minimal model for interchange motions describes the coupling between a thermodynamic variable and the plasma vorticity. The analysis focuses on classification of topological changes of the structures as they develop in time and as the Rayleigh number varies.**

## INTRODUCTION

At the edge region of a magnetically confined toroidal plasma the transport of particles and energy is dominated by recurring outbreaks of coherent plasma structures called blobs<sup>1</sup>. In fluid dynamics similarly shaped structures known as plumes<sup>2</sup> arise when a fluid in a gravitational field is driven by thermal convection. This work focuses on the evolution of the blob topology during propagation.

## MINIMAL MODEL FOR INTERCHANGE MOTIONS

We consider a minimal interchange model<sup>1</sup>:

$$\begin{aligned} \left( \frac{\partial}{\partial t} + \hat{\mathbf{z}} \times \nabla \phi \cdot \nabla \right) \theta &= \kappa \nabla_{\perp}^2 \theta, \\ \left( \frac{\partial}{\partial t} + \hat{\mathbf{z}} \times \nabla \phi \cdot \nabla \right) \Omega + \frac{\partial \theta}{\partial y} &= \nu \nabla_{\perp}^2 \Omega, \end{aligned}$$

and  $\nabla_{\perp}^2 \phi = \Omega$ . Here  $\theta(x, y, t)$  is the thermodynamic variable (e.g. density, pressure or temperature),  $\Omega(x, y, t)$  the vorticity, and  $\phi(x, y, t)$  the electrostatic potential.  $\kappa$  and  $\nu$  are parameters related to the Rayleigh and Prandtl numbers by  $Ra = 1/(\kappa\nu)$  and  $Pr = \nu/\kappa$ .  $x \in [-20, 30]$  and  $y \in [-25, 25]$  are the spatial coordinates and  $t \in [0, 20]$  is time. Dirichlet boundary conditions were applied on all boundaries.

All simulations were initialized as

$$\begin{aligned} \theta(x, y, t=0) &= e^{-\frac{1}{2}(x^2+y^2)}, \\ \Omega(x, y, t=0) &= 0, \\ \phi(x, y, t=0) &= 0. \end{aligned}$$




Figure: Plot of  $\theta(x, y, t=0)$  and the two initial nullclines.

## A DYNAMICAL SYSTEMS APPROACH

Streamlines of the thermodynamic variable  $\theta$  are determined from

$$\begin{pmatrix} \partial_s x \\ \partial_s y \end{pmatrix} = \begin{pmatrix} -\partial_y \theta(x, y, t_0; Ra) \\ \partial_x \theta(x, y, t_0; Ra) \end{pmatrix} \quad (1)$$

Since  $d\theta/ds = 0$  these streamlines are level curves of  $\theta$ . We shall investigate bifurcations of the streamline topology as  $t$  and  $Ra$  varies. Equilibrium points of the system (1) are stationary points of  $\theta$ . Intersections of the nullclines of the system (1) determines the stationary points. The eigenvalues of the Hessian of  $\theta$  evaluated at a stationary point determines the type (local maximum, local minimum, or saddle) of the stationary point [3].

## BIFURCATION ANALYSIS




Figure: Bifurcation diagram with  $t$  and  $Ra$  as bifurcation parameters. “PF” indicates a pitchfork bifurcation and “SC” a saddle-center bifurcation. Solid and dashed lines indicate bifurcations involving local maxima and local minima, respectively.

## THE DIFFERENT STRUCTURES

The bifurcation curves in the bifurcation diagram separates the  $(t, Ra)$ -parameter space into 18 regions. Within each region any set of streamline patterns of  $\theta$  are topologically equivalent.




Figure: A representative solution  $\theta(x, y, t_0)$  for each of the 18 regions in the bifurcation diagram. Black dots indicate local maxima, white dots indicate local minima and crosses indicate saddles. Local maxima and minima of  $\theta$  correspond to centers of the streamlines of  $\theta$  and saddles of  $\theta$  correspond to topological saddles of the streamlines of  $\theta$ .

## CONCLUSION AND OUTLOOK

We have carried out a bifurcation analysis of the topological changes of seeded blobs using time and Rayleigh number as parameters. A larger Rayleigh number and time give rise to a more complicated topology with more stationary points. Future work includes:

- Investigate dependence on the Prandtl number.
- Compare with a similar bifurcation diagram for the vorticity.
- Determine measurable characteristics of the different topological structures such as propagation speed.

## References:

- <sup>1</sup>O. E. Garcia, N. H. Bian, and W. Fundamenski, Phys. Plasmas **13**, 082309 (2006).
- <sup>2</sup>T. Bohr, M. Jensen, G. Paladin, and A. Vulpiani, *Dynamical systems approach to turbulence*, (Cambridge University Press, 1998).
- <sup>3</sup>M. Brøns, Adv. Appl. Mech. **41**, 1–42 (2007).